



Lower Colorado River Multi-Species Conservation Program

Balancing Resource Use and Conservation

Post-Development and System-Wide Monitoring of Rodent Populations

2019 Annual Report



March 2021

Work conducted under LCR MSCP Work Tasks D10 and F3

Lower Colorado River Multi-Species Conservation Program

Steering Committee Members

Federal Participant Group

Bureau of Reclamation
U.S. Fish and Wildlife Service
National Park Service
Bureau of Land Management
Bureau of Indian Affairs
Western Area Power Administration

Arizona Participant Group

Arizona Department of Water Resources
Arizona Electric Power Cooperative, Inc.
Arizona Game and Fish Department
Arizona Power Authority
Central Arizona Water Conservation District
Cibola Valley Irrigation and Drainage District
City of Bullhead City
City of Lake Havasu City
City of Mesa
City of Somerton
City of Yuma
Electrical District No. 3, Pinal County, Arizona
Golden Shores Water Conservation District
Mohave County Water Authority
Mohave Valley Irrigation and Drainage District
Mohave Water Conservation District
North Gila Valley Irrigation and Drainage District
Town of Fredonia
Town of Thatcher
Town of Wickenburg
Salt River Project Agricultural Improvement and Power District
Unit "B" Irrigation and Drainage District
Wellton-Mohawk Irrigation and Drainage District
Yuma County Water Users' Association
Yuma Irrigation District
Yuma Mesa Irrigation and Drainage District

Other Interested Parties Participant Group

QuadState Local Governments Authority
Desert Wildlife Unlimited

California Participant Group

California Department of Fish and Wildlife
City of Needles
Coachella Valley Water District
Colorado River Board of California
Bard Water District
Imperial Irrigation District
Los Angeles Department of Water and Power
Palo Verde Irrigation District
San Diego County Water Authority
Southern California Edison Company
Southern California Public Power Authority
The Metropolitan Water District of Southern California

Nevada Participant Group


Colorado River Commission of Nevada
Nevada Department of Wildlife
Southern Nevada Water Authority
Colorado River Commission Power Users
Basic Water Company

Native American Participant Group

Hualapai Tribe
Colorado River Indian Tribes
Chemehuevi Indian Tribe

Conservation Participant Group

Ducks Unlimited
Lower Colorado River RC&D Area, Inc.
The Nature Conservancy



— BUREAU OF —
RECLAMATION

Lower Colorado River Multi-Species Conservation Program

Post-Development and System-Wide Monitoring of Rodent Populations

2019 Annual Report

Prepared by:

Jeff Hill and Joe Lyon, Wildlife Group

**Lower Colorado River
Multi-Species Conservation Program
Bureau of Reclamation
Lower Colorado Basin
Boulder City, Nevada
<http://www.lcrmscp.gov>**

March 2021

Hill, J. and J. Lyon. 2021. Post-Development and System-Wide Monitoring of Rodent Populations, 2019 Annual Report. Lower Colorado River Multi-Species Conservation Program, Boulder City, Nevada.

ACRONYMS AND ABBREVIATIONS

Cibola NWR	Cibola National Wildlife Refuge
km	kilometer(s)
LCR	lower Colorado River
LCR MSCP	Lower Colorado River Multi-Species Conservation Program
Nature Trail	Cibola National Wildlife Refuge Unit #1 Conservation Area Nature Trail

CONTENTS

	Page
Abstract.....	iii
Introduction.....	1
Methods.....	2
Study Areas.....	3
LCR MSCP Conservation Areas	5
Beal Lake Conservation Area	5
Palo Verde Ecological Preserve.....	7
Cibola National Wildlife Refuge Unit #1 Conservation Area	9
Cibola Valley Conservation Area	11
Hart Mine Marsh.....	13
Laguna Division Conservation Area.....	15
Yuma East Wetlands.....	17
Hunters Hole	19
Results	21
Conservation Areas	21
Beal Lake Conservation Area	21
Palo Verde Ecological Reserve.....	23
Cibola National Wildlife Refuge Unit #1 Conservation Area	25
Cibola Valley Conservation Area	27
Hart Mine Marsh.....	28
Laguna Division Conservation Area.....	29
Yuma East Wetlands.....	30
Hunters Hole	32
Literature Cited	35

Tables

Table	Page
1 Beal Lake Conservation Area captures.....	22
2 Palo Verde Ecological Reserve Phase 4 captures.....	24
3 Palo Verde Ecological Reserve Phase 8 captures.....	24
4 Cibola National Wildlife Refuge Unit #1 Conservation Area Hippy Fire captures.....	26
5 Cibola National Wildlife Refuge Unit #1 Conservation Area Nature Trail captures.....	26
6 Cibola Valley Conservation Area captures.....	27
7 Hart Mine Marsh captures.....	28
8 Laguna Division Conservation Area captures.....	29
9 Yuma East Wetlands Zone C captures.....	31
10 Yuma East Wetlands Zone F captures.....	31
11 Hunters Hole captures.....	33

Figures

Figure	Page
1 LCR MSCP conservation areas.....	4
2 Beal Lake Conservation Area managed acreage through fiscal year 2019.....	6
3 Palo Verde Ecological Reserve managed acreage through fiscal year 2019.....	8
4 Cibola National Wildlife Refuge Unit #1 Conservation Area managed acreage through fiscal year 2019.....	10
5 Cibola Valley Conservation Area managed acreage through fiscal year 2019.....	12
6 Hart Mine Marsh managed acreage through fiscal year 2019.....	14
7 Laguna Division Conservation Area managed acreage through fiscal year 2019.....	16
8 Yuma East Wetlands managed acreage through fiscal year 2019.....	18
9 Hunters Hole managed acreage through fiscal year 2019.....	20
10 Beal Lake Conservation Area trapping areas.....	21
11 Palo Verde Ecological Reserve trapping areas.....	23
12 Cibola National Wildlife Refuge Unit #1 Conservation Area trapping areas.....	25
13 Cibola Valley Conservation Area trapping areas.....	27
14 Hart Mine Marsh trapping areas.....	28
15 Laguna Division Conservation Area survey areas.....	29
16 Yuma East Wetlands trapping areas.....	30
17 Hunters Hole trapping areas.....	32

ABSTRACT

The Bureau of Reclamation is the lead agency for the Lower Colorado River Multi-Species Conservation Program (LCR MSCP). Implementing the LCR MSCP will create at least 8,132 acres of new habitat (5,940 acres of Fremont cottonwood-Goodding's willow [*Populus fremontii*-*Salix gooddingii*], 1,320 acres of honey mesquite [*Prosopis glandulosa*], 512 acres of marsh, and 360 acres of backwater) for 27 covered species, including the Colorado River cotton rat (*Sigmodon arizonae plenus*) and Yuma hispid cotton rat (*Sigmodon hispidus eremicus*). The *sobrinus* subspecies of the desert pocket mouse (*Chaetodipus penicillatus*) is being evaluated to determine if it should be added as a covered species.

The presence of Colorado River cotton rats, Yuma hispid cotton rats, and desert pocket mice at existing habitat along the lower Colorado River and at LCR MSCP conservation areas was monitored under Work Task D10 (System Monitoring of Rodent Populations) and Work Task F3 (Small Mammal Colonialization of Conservation Areas). Trapping was conducted in eight conservation areas during fiscal year 2019. Eighteen Colorado River cotton rat captures were recorded at the following conservation areas: Cibola Valley Conservation Area, Palo Verde Ecological Reserve, and Hart Mine Marsh. Six Yuma hispid cotton rats were captured at the following conservation areas: Laguna Division Conservation Area and Hunters Hole. Sixteen desert pocket mice were captured in the following conservation areas: Cibola Valley Conservation Area, Yuma East Wetlands, Hunters Hole, Palo Verde Ecological Reserve, and the Laguna Division Conservation Area.

A genetic analysis was not conducted on desert pocket mice, so it is unknown if they were of the *sobrinus* subspecies, which is thought to be restricted to southeast Nevada, northwestern Arizona, and extreme southwestern Utah.

INTRODUCTION

The Lower Colorado River Multi-Species Conservation Program (LCR MSCP) is a partnership of Federal and non-Federal stakeholders that was created to respond to the need to balance the use of lower Colorado River (LCR) water resources and the conservation of native species and their habitats in compliance with the Endangered Species Act. It is a long-term (50-year) plan to conserve at least 27 species along the LCR from Lake Mead to the Southerly International Boundary with Mexico through implementation of a Habitat Conservation Plan. Implementation of the LCR MSCP began in October 2005. Implementing the LCR MSCP will create at least 8,132 acres of new habitat (5,940 acres of Fremont cottonwood-Goodding's willow [*Populus fremontii-Salix gooddingii*] [hereafter cottonwood-willow], 1,320 acres of honey mesquite [*Prosopis glandulosa*], 512 acres of marsh, and 360 acres of backwater).

Twenty-seven Federal or State-listed candidate and sensitive species and their associated habitats, ranging from aquatic and wetland habitats to riparian and upland areas, are covered under the LCR MSCP. These include the Colorado River cotton rat (*Sigmodon arizonae plenus*) and Yuma hispid cotton rat (*Sigmodon hispidus eremicus*). The *sobrinus* subspecies of the desert pocket mouse (*Chaetodipus penicillatus*) is being evaluated to determine if it should be added as a covered species.

Within the LCR MSCP habitat creation goals, 125 acres of habitat are to be created for the Colorado River cotton rat, and 76 acres are to be created for the Yuma hispid cotton rat. It is currently believed that the ranges of these two cotton rat species do not overlap. Those captured south of the Trigo and Chocolate Mountains in the area of the Imperial National Wildlife Refuge and south of the Yuma, Arizona, area to date are Yuma hispid cotton rats. Those captured north of the aforementioned mountain ranges to date are Colorado River cotton rats. The northernmost historic records of Colorado River cotton rats are from an area just south of Laughlin, Nevada (Bradley 1966; Hall 1946). Colorado River cotton rats were captured in 2012 at the Big Bend Conservation Area, marking the first record of the species in Nevada since 1966.

Desert pocket mice occur in creosote bush (*Larrea tridentata*) and xeric riparian communities of the Southwest, from Baja California, Mexico in the South and southeastern California, southern Nevada, and extreme southwestern Utah in the North. The range of the *sobrinus* subspecies is not well documented, but it is believed to be within Clark County, Nevada, and may be present farther south. The desert pocket mice caught during the 2017 small mammal surveys were not evaluated to determine whether or not they were of the *sobrinus* subspecies.

Post-Development and System-Wide Monitoring of Rodent Populations 2019 Annual Report

The Bureau of Reclamation is increasing its understanding of restoration science through an adaptive management approach; therefore, monitoring of habitat creation/restoration sites is crucial. Species presence at existing habitat along the LCR and at LCR MSCP conservation areas is being monitored under Work Task D10 (System Monitoring of Rodent Populations) and Work Task F3 (Small Mammal Colonialization of Conservation Areas). In addition, Work Task C27 (Small Mammal Population Studies) was conducted to identify distribution, genetics, and habitat requirements, and to establish monitoring protocols for the covered small mammal species.

METHODS

Sherman live traps (3 x 3.75 x 12 inches) were placed non-randomly in areas with the highest likelihood of encountering the cotton rat species, which generally consist of a dense understory of grasses and/or shrubs alongside or beneath an overhead cover of forbs and low-growing woody species (Work Task C27) (Goertz 1964). In general, if the overall trapping area is fairly homogenous, with appropriately dense vegetation structure, the trapping grid begins on a corner or edge. If a distinctly denser vegetation area is found within an otherwise more open area, the trapping grid begins in the denser habitat patch. These surveys were focused on determining the presence of the cotton rat species. All other captures were incidental to the main focus, and the methods were not designed to estimate abundance; therefore, the numbers of individuals of each species (including Colorado River cotton rats and Yuma hispid cotton rats) reported here are rough indices that do not provide reliable means of analyzing population trends.

Sherman live traps are triggered by an animal stepping on a pressure plate that then closes a trap door behind it. Traps were baited with a mixture of oats, peanut butter, and vanilla. A small handful of cotton was also added to each trap to provide insulating cover for any animal trapped overnight. The traps were set 1 to 2 hours before sunset and pulled within 1 to 2 hours of sunrise during the warmer months to minimize heating of the traps during the trapping period.

When the habitat patch allowed, traps were set out in a grid of at least three transects approximately 50 feet (15 meters) apart. Each transect consisted of at least five trap stations spaced approximately 33 feet (10 meters) apart. Trap spacing is estimated with pacing and is not measured exactly; it may vary if the more densely vegetated areas are dispersed across the trapping area.

When the habitat patch with the highest density vegetation was more linear, transects were not set in a grid system. If only a single linear stretch of habitat was trapped, a single transect was set along the edge of the habitat patch. This usually meant that the transect was much longer than in the typical grid system (15 or more trap stations).

The approximate location of each trap was recorded using an Apple iPhone 6 or iPad Mini 4 with ArcGIS Survey123 software installed, and the traps were collected the following morning. Each transect was labeled by a letter, and each trap was numbered. For example, the first trap station of the first transect of a grid would be labeled A-1, and if there were two traps at that location, they were designated as trap A-1-a and A-1-b. Data were entered electronically on the Global Positioning System unit using a mobile electronic field form.

Captured animals were either viewed inside the trap or transferred into a clear plastic bag and identified to species. The animals were identified using the Mammals of California field guide (Jameson and Peeters 2004), the Kays and Wilson (2002) field guide, and the expertise of LCR MSCP personnel that had more than 3 years of experience conducting surveys. Cotton rats were given an age class based on three general size classes (juvenile, subadult, adult). The age classes of the cotton rats were determined based on the general size of an individual using the experience of LCR MSCP personnel present during the survey. The gender of the cotton rats was also determined. All other species were only given an age class of either juvenile or adult, and gender was not determined. Measurements were taken, if needed, for identification. A mobile electronic field form was used to record information on all animals captured, whether they were marked, where in the grid they were captured, the location of the grid, and what the dominant ground cover vegetation was in the trapping area. All animals were released back into the trapping area once identification was made. Traps in which an animal had been captured were washed in a bleach water solution, rinsed in plain water, and then set out to dry after each trapping day.

STUDY AREAS

Trapping was conducted in seven LCR MSCP conservation areas within Reaches 2–7 (figure 1).

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**




Figure 1.—LCR MSCP conservation areas.

LCR MSCP Conservation Areas

Beal Lake Conservation Area

The Beal Lake Conservation Area is 100 acres adjacent to Beal Lake and Topock Marsh, inside the Havasu National Wildlife Refuge on the Arizona side of the Colorado River (see figure 1). The site is within Reach 3. The site contains 119 acres of cottonwood-willow habitat and 225 acres backwater habitat (figure 2). It is a two-phase habitat creation project that was initiated in spring 2003. The site was planted with Fremont cottonwood, Goodding's willow, coyote willow (*Salix exigua*), honey mesquite, and screwbean mesquite (*Prosopis pubescens*). Arrowweed (*Pluchea sericea*) and some baccharis (*Baccharis* spp.) have begun to fill in the open areas and edges of most of the plots in the site.

Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report




Figure 2.—Beal Lake Conservation Area managed acreage through fiscal year 2019.

Palo Verde Ecological Preserve

The Palo Verde Ecological Preserve is a conservation area located 5 miles (8 kilometers [km]) north of Blythe, California, along the California side of the Colorado River (see figure 1). The site is within Reach 4. It encompasses 1,300 acres. The acreages are separated into nine different phases, with one phase planted every year through 2014. There are currently 950 cottonwood-willow acres planted and 78 acres of honey mesquite (figure 3). In spring 2006, a 31-acre nursery (Phase 1) was planted. In spring 2007, Phase 2 was planted with 80 acres of Fremont cottonwood, Goodding's willow, coyote willow, and other riparian plants. Phase 3 was planted in spring 2008 and is also planted with cottonwood-willow habitat types. Phase 4 was planted in 2009 and contains mostly cottonwood-willow, with one plot of honey mesquite and a mix of native grasses. Phases 5, 6, and 7 were planted in 2010, 2011, and 2012, respectively, with cottonwood-willow habitat. Phase 8 was planted in 2013 with 38 acres of honey mesquite and quailbush (*Atriplex lentiformis*) habitat.

Post-Development and System-Wide Monitoring of Rodent Populations
 2019 Annual Report




Figure 3.—Palo Verde Ecological Reserve managed acreage through fiscal year 2019.

Cibola National Wildlife Refuge Unit #1 Conservation Area

The Cibola National Wildlife Refuge (Cibola NWR) is located approximately 30 miles (48 km) south of Blythe, California, along 12 miles (19 km) of the LCR in Arizona and California (see figure 1). The site is within Reach 4. The Cibola NWR is divided into six management units, of which the Cibola National Wildlife Refuge Unit #1 Conservation Area comprises approximately 949 acres (figure 4). The Lower Hippy Fire area was planted in fiscal year 2016 with 96 acres of cottonwood-willow habitat. There are emergent grasses throughout the area, as the cottonwood canopy cover has not shaded the ground out yet.

Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report




Figure 4.—Cibola National Wildlife Refuge Unit #1 Conservation Area managed acreage through fiscal year 2019.

Cibola Valley Conservation Area

The Cibola Valley Conservation Area is located in Arizona adjacent to the Colorado River, about 15 miles (24 km) south of Blythe, California (see figure 1). The site is within Reach 4. It will encompass about 1,235 acres when completed. Three phases included the planting of Fremont cottonwood, Goodding's willow, coyote willow, and other riparian plant species. Phase 1 was planted in spring 2006 and contains a 22-acre nursery and a 64-acre area of cottonwood-willow habitat. Phase 3 was planted in spring 2007 and contains over 80 acres of cottonwood-willow planted in different combinations (figure 5). Phase 3 also includes 11 acres of baccharis mixed with some cottonwood-willow. Phase 2 was planted in spring 2008. Most of Phase 2 is planted with cottonwood-willow habitat, with one small area of honey mesquite and quailbush. Phase 4 was planted in 2009 with honey mesquite and quailbush. Phase 5 was planted in 2010 with 71 acres of honey mesquite and quailbush. Phase 6 was planted in 2011 with 89 acres of honey mesquite and quailbush.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**




Figure 5.—Cibola Valley Conservation Area managed acreage through fiscal year 2019.

Hart Mine Marsh

Hart Mine Marsh is located on the southern part of the U.S. Fish and Wildlife Service's Cibola NWR in Arizona (figure 6). It makes up the lower part of the management area defined as Unit 2 by the Cibola NWR. Approximately 255 acres of the 893-acre site have been restored and are being managed to partially fulfill the marsh habitat creation goals of the LCR MSCP and have been designated as a LCR MSCP conservation area. The remaining 515 acres with wetland potential are managed for U.S. Fish and Wildlife Service specific goals but are intended to be additive to the wetland complex's habitat mosaic.

Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report




Figure 6.—Hart Mine Marsh managed acreage through fiscal year 2019.

Laguna Division Conservation Area

The Laguna Division Conservation Area is located 20 miles north of Yuma, Arizona, between Imperial Dam and Laguna Dam (figure 7). The site is within Reach 6. It covers 1,200 acres, straddling the California/Arizona State boundary. The Laguna Division Conservation Area comprises 1,129 acres of cottonwood-willow and 42 acres of honey mesquite habitat. Planting was completed in 2016.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**




Figure 7.—Laguna Division Conservation Area managed acreage through fiscal year 2019.

Yuma East Wetlands

Yuma East Wetlands is a 392-acre conservation area consisting of lands owned by the Quechan Indian Tribe, city of Yuma, and the Arizona Game and Fish Department. The site is within Reach 6. The site is located within the city of Yuma, Arizona (see figure 1). Habitat restoration activities began in Yuma East Wetlands in 2004. The majority of the planting took place in 2009, including 183 acres of cottonwood-willow habitat, 94 acres of marsh habitat, and 103 acres of honey mesquite habitat (figure 8). In 2013, the LCR MSCP entered into partnership with the Quechan Indian Tribe, city of Yuma, Arizona Game and Fish Commission, and the Yuma Crossing National Heritage Area to support the long-term management of the site. In 2014, the bankline portion of the site was cleared of invasive vegetation and replanted with native vegetation. The site contains many open canopy areas that allow for a healthy variety of shrub and grass species necessary for Yuma hispid cotton rat colonization. In 2018, a large fire swept through a portion of the cottonwood-willow habitat, clearing much of the grasses inhabited by Yuma hispid cotton rats.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**



Figure 8.—Yuma East Wetlands managed acreage through fiscal year 2019.

Hunters Hole

Hunters Hole is a conservation area located near the community of San Luis, 18 miles (29 km) south of Yuma, Arizona (see figure 1). The site is within Reach 7. It was added to the LCR MSCP in 2011. The site was planted in spring 2012 with approximately 44 acres of cottonwood-willow habitat (figure 9) consisting of cottonwood-willow, honey mesquite, and marsh. A number of fires have occurred at Hunters Hole throughout the years, which periodically open up the habitat for a dense layer of grasses to grow in before being shaded out by canopy regrowth.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**




Figure 9.—Hunters Hole managed acreage through fiscal year 2019.

RESULTS

Conservation Areas

Beal Lake Conservation Area

Trapping occurred in planted cottonwood-willow habitat within the Beal Lake Conservation Area. The traps were set within a narrow patch of arrowweed and various grass species that have potentially suitable structure for Colorado River cotton rats (figure 10). In recent years, the patch has narrowed, becoming increasingly dominated by arrowweed. Fall 2018 trapping resulted in no captures (table 1). Two traps were removed on the second night of trapping due to the presence of ants. No trapping was conducted in spring 2019.



Figure 10.—Beal Lake Conservation Area trapping areas.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Table 1.—Beal Lake Conservation Area captures

Species	Fall 2018 (November 27–28, 2018)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	0
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	0
Cactus mouse (<i>Peromyscus eremicus</i>)	0
White-throated wood rat (<i>Neotoma albigula</i>)	0
Total captures	0
Traps/nights	40/1, 38/1

Palo Verde Ecological Reserve

This site consists of multiple planting phases with a variety of habitat types. All of the areas trapped contained a grass and/or shrub component. Phases 4 and 8 were trapped. Phase 4 was planted with honey mesquite and quailbush and is now intermixed with Bermudagrass (*Cynodon dactylon*) and other grasses and forbs. Phase 8 was planted with honey mesquite in 2013. Trapping occurred in Phases 4 and 8 (figure 11) over 2 nights. No trapping was conducted in fall 2018. Spring 2019 trapping sessions produced one desert pocket mouse and one Colorado River cotton rat capture; a desert pocket mouse was encountered in Phase 4 (table 2), and a Colorado River cotton rat was encountered in Phase 8 (table 3).




Figure 11.—Palo Verde Ecological Reserve trapping areas.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Table 2.—Palo Verde Ecological Reserve Phase 4 captures

Species	Spring 2019 (February 26–27, 2019)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	0
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	1
Cactus mouse (<i>Peromyscus eremicus</i>)	9
White-throated wood rat (<i>Neotoma albigula</i>)	1
Total captures	11
Traps/nights	40/2

Table 3.—Palo Verde Ecological Reserve Phase 8 captures

Species	Spring 2019 (February 26–27, 2019)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	1
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	0
Cactus mouse (<i>Peromyscus eremicus</i>)	16
Deer mouse (<i>Peromyscus maniculatus</i>)	4
Harvest mouse (<i>Reithrodontomys megalotis</i>)	1
Total captures	22
Traps/nights	40/2

Cibola National Wildlife Refuge Unit #1 Conservation Area

Surveys within the Cibola National Wildlife Refuge Unit #1 Conservation Area occurred in the Lower Hippy Fire section and the Cibola National Wildlife Refuge Unit #1 Conservation Area Nature Trail (Nature Trail). The Hippy Fire area consists of young cottonwoods and emergent Bermudagrass (figure 12). Trapping in the Nature Trail occurred in an area in the middle of the site dominated by Johnsongrass (*Sorghum halepense*) and baccharis and at an area on the northern part of the site dominated by Bermudagrass and cottonwoods. No covered species were captured over 2 nights of fall trapping at the Hippy Fire area (table 4). Four Colorado River cotton rats and no desert pocket mice were captured over 2 nights of trapping in fall 2018 in the Nature Trail (table 5). No traps were set during spring 2019.




Figure 12.—Cibola National Wildlife Refuge Unit #1 Conservation Area trapping areas.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Table 4.—Cibola National Wildlife Refuge Unit #1 Conservation Area Hippy Fire captures

Species	Fall 2018 (November 29–30, 2018)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	0
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	0
Cactus mouse (<i>Peromyscus eremicus</i>)	2
Deer mouse (<i>Peromyscus maniculatus</i>)	5
Harvest mouse (<i>Reithrodontomys megalotis</i>)	10
White-throated wood rat (<i>Neotoma albigula</i>)	0
Total captures	17
Traps/nights	80/2

Table 5.—Cibola National Wildlife Refuge Unit #1 Conservation Area Nature Trail captures

Species	Fall 2018 (November 29–30, 2018)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	4
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	0
White-throated wood rat (<i>Neotoma albigula</i>)	4
Cactus mouse (<i>Peromyscus eremicus</i>)	12
Harvest mouse (<i>Reithrodontomys megalotis</i>)	1
Deer mouse (<i>Peromyscus maniculatus</i>)	1
Total captures	22
Traps/nights	40/2

Cibola Valley Conservation Area

Phase 1 of the Cibola Valley Conservation Area was trapped for 2 nights in fall 2018 (figure 13). One trapping area was along a strip of mule-fat (*Baccharis salicifolia*), where the shrubs met a dense area of Bermudagrass. The other trapping area consisted of dead coyote willow and Bermudagrass. Six Colorado River cotton rats were captured (table 6). No traps were set during spring 2019




Figure 13.—Cibola Valley Conservation Area trapping areas.

Table 6.—Cibola Valley Conservation Area captures

Species	Fall 2018 (November 27–28 2018)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	6
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	0
Cactus mouse (<i>Peromyscus eremicus</i>)	10
Deer mouse (<i>Peromyscus maniculatus</i>)	6
Harvest mouse (<i>Reithrodontomys megalotis</i>)	14
Total captures	36
Traps/nights	160/2

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Hart Mine Marsh

Surveys at Hart Mine Marsh occurred in Cells 01 and 02, in three trapping grids set up in between arrowweed and some cattails (*Typha* spp.)/emergent grasses at the water’s edge (figure 14). No surveys were conducted during fall 2018. Seven Colorado River cotton rats and one desert pocket mouse were captured over 2 nights of trapping in spring 2019 (table 7).




Figure 14.—Hart Mine Marsh trapping areas.

Table 7.—Hart Mine Marsh captures

Species	Spring 2019 (February 28 – March 1, 2019)
Colorado River cotton rat (<i>Sigmodon arizonae plenus</i>)	7
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	1
Cactus mouse (<i>Peromyscus eremicus</i>)	3
Harvest mouse (<i>Reithrodontomys megalotis</i>)	7
Total captures	18
Traps/nights	120/2

Laguna Division Conservation Area

Trapping occurred in three transects within the Laguna Division Conservation Area. Trapping areas were comprised of sparse alkali sacaton (*Sporobolus airoides*) and arrowweed (figure 15). Two nights of trapping in fall 2018 resulted in the capture of 5 Yuma hispid cotton rats and 12 desert pocket mice (table 8). No surveys were conducted during spring 2019.




Figure 15.—Laguna Division Conservation Area survey areas.

Table 8.—Laguna Division Conservation Area captures

Species	Fall 2018 (October 18–19, 2018)
Yuma hispid cotton rat (<i>Sigmodon hispidus eremicus</i>)	5
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	12
Cactus mouse (<i>Peromyscus eremicus</i>)	12
House mouse (<i>Mus musculus</i>)	2
Merriam's kangaroo rat (<i>Dipodomys merriami</i>)	2
Total captures	33
Traps/nights	180/2

Yuma East Wetlands

Surveys occurred in Zones C and F. The trapping area consisted of a mix of alkali sacaton, baccharis, and some honey mesquite plantings (figure 16). Trapping in Zone C was focused along a previous burned area, where new grass is growing in, and in an unburned mix of baccharis and honey mesquite. No surveys were conducted in fall 2018. Two nights of surveys in spring 2019 encountered no Yuma hispid cotton rats. No LCR MSCP species were captured in Zone C (table 9). One desert pocket mouse was captured in Zone F (table 10).




Figure 16.—Yuma East Wetlands trapping areas.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Table 9.—Yuma East Wetlands Zone C captures

Species	Spring 2019 (March 5–6, 2019)
Yuma hispid cotton rat (<i>Sigmodon hispidus eremicus</i>)	0
Cactus mouse (<i>Peromyscus eremicus</i>)	13
Harvest mouse (<i>Reithrodontomys megalotis</i>)	1
Total captures	14
Traps/nights	40/2

Table 10.—Yuma East Wetlands Zone F captures

Species	Spring 2019 (March 5–6, 2019)
Yuma hispid cotton rat (<i>Sigmodon hispidus eremicus</i>)	0
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	1
Cactus mouse (<i>Peromyscus eremicus</i>)	10
Harvest mouse (<i>Reithrodontomys megalotis</i>)	1
Total captures	12
Traps/nights	40/2

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Hunters Hole

Surveys occurred in Cells 02 and 03 over 2 nights each in fall 2018 and spring 2019. The trapping areas were planted with Fremont cottonwood, Goodding’s willow, coyote willow, and honey mesquite along with alkali sacaton and blue grama (*Bouteloua gracilis*) (figure 17). Both trapping areas were dominated by grasses, though trees are beginning to shade out the ground cover. Fall 2018 surveys produced no Yuma hispid cotton rats or desert pocket mice (table 11). Spring 2019 surveys produced one Yuma hispid cotton rat and one desert pocket mouse (table 11).



Figure 17.—Hunters Hole trapping areas.

**Post-Development and System-Wide Monitoring of Rodent Populations
2019 Annual Report**

Table 11.—Hunters Hole captures

Species	Fall 2018 (October 16–17, 2018)	Spring 2019 (March 7–8, 2019)
Yuma hispid cotton rat (<i>Sigmodon hispidus eremicus</i>)	0	1
Desert pocket mouse (<i>Chaetodipus penicillatus</i> sp.)	0	1
Cactus mouse (<i>Peromyscus eremicus</i>)	10	13
Deer mouse (<i>Peromyscus maniculatus</i>)	2	0
White-throated wood rat (<i>Neotoma albigula</i>)	0	1
Total captures	12	16
Traps/nights	60/1, 80/1	80/2

LITERATURE CITED

- Bradley, W.G. 1966. Status of the cotton rat in Nevada. *Journal of Mammalogy* 47:349–350.
- Goertz, J.W. 1964. The influence of habitat quality upon density of cotton rat populations. *Ecological Monographs*:359–381.
- Hall, E.R. 1946. *Mammals of Nevada*. University of Nevada Press, Reno and Las Vegas, Nevada.
- Jameson, E.W. and H.J. Peeters. 2004. *Mammals of California; California Natural History Guide No. 66*. University of California Press, Berkeley and Los Angeles, California.
- Kays, R.W. and D.E. Wilson. 2002. *Mammals of North America*. Princeton University Press, Princeton, New Jersey.